

# Preventive Care

## Stay Healthy, Save Money

Preventing and detecting disease early is important to living a healthy life. The better your health, the lower your health care costs are likely to be. One of the most important actions you can take for your health and your family's health is to schedule regular check-ups and screenings with your primary care physician.

Your State of Ohio medical plan – Ohio Med PPO – offers the following services with no deductible, no copayment and no coinsurance for network providers. Other services are available for the normal copayment, coinsurance and deductible amounts.

FREE EXAMS AND SCREENINGS	
Clinical breast exam	1/plan year
Colonoscopy	Every 10 years starting at age 50
Flexible sigmoidoscopy	Every 10 years starting at age 50
Glucose	1/plan year
Gynecological Exam	1/plan year
Hemoglobin, hematocrit or CBC	1/plan year
Lipid profile or total and HDL cholesterol	1/plan year
Mammogram	1 routine and 1 medically necessary/plan year
Pre-natal office visits	As needed; based on physician's ability to code claims separately from other maternity-related services
Prostate-specific Antigen (PSA)	1/plan year starting at age 40
Stool for occult blood	1/plan year
Urinalysis	1/plan year
Well-baby, well-child exam	Various for birth to 2 years; then annual to age 21
Well-person exam (annual physical)	1/plan year

FREE IMMUNIZATIONS	
Diphtheria, tetanus, pertussis (DTap)	2, 4, 6 and 15-18 months; 4-6 years
Haemophilus influenza b (Hib)	2, 4, 6 and 12-15 months
Hepatitis A (HepA)	2 doses between 1-2 years
Hepatitis B (HepB)	Birth; 1-2 months; 6-18 months
Human Papillomavirus (HPV)	3 doses for 9-26 years
Influenza	1/plan year
Measles, mumps, rubella (MMR)	12-15 months, then at 4-6 years; adults who lack immunity
Meningococcal (MCV4)	1 dose between 11-12 years or start of high school or college
Pneumococcal	2, 4 and 6 months; 12-15 months; annually at age 65 and older; high risk groups
Poliovirus (IPEV)	2 and 4 months; 6-18 months; 4-6 years
Rotavirus (Rota)	2, 4 and 6 months
Tetanus, diphtheria, pertussis (Td/Tdap)	11-12 years; Td booster every 10 years, 18 and older
Varicella (Chickenpox)	12-15 months; 4-6 years; 2 doses for susceptible adults
Zoster (shingles)	1 dose for age 19 +

Note: This is not an all-inclusive list. Please refer to [das.ohio.gov/medical](https://das.ohio.gov/medical) for more information about preventive care services.